



## Goat Anti-Mouse IgG: DyLight® 488


Cat No: 16700 - 1 mg

### General Data

**Shipping:** wet ice

**Formulation:** A liquid

**Isotype:** IgG



Detection of mouse IgG using Columbia Biosciences anti-mouse IgG conjugated to DyLight® 488: Raji B cells were stained for 30 minutes with 5 µg/ml Columbia Biosciences anti-mouse IgG conjugated to DyLight 488 after treatment for 1 hour with staining buffer alone (white histogram), staining buffer containing mouse anti-human CD45 primary antibody (5 µg/ml; black histogram), or staining buffer containing mouse IgG isotype control (red-outlined histogram). 10,000 cells per sample were analyzed on a flow cytometer for fluorescence emission at 530 nm.

**Application(s):** FC, FACS, Protein Microarrays

### Product Overview

Goat anti-Mouse IgG, secondary antibody, conjugated to DyLight® 488 Antibody: Goat polyclonal anti-Mouse IgG Dye: DyLight® 488 Excitation max. λ: 493 nm Emission max. λ: 518 nm Uses: Flow cytometry, cell-based assays, microarrays, and microplate applications Form: Conjugate supplied as a liquid in 0.1 M sodium phosphate, 0.15 M NaCl (pH7.4), 1% BSA, and 0.02% NaN<sub>3</sub>.

FP/22/24

For research laboratory use only – Not for human diagnostic use.

Buyers agree to purchase the material subject to Purchasing Terms that can be found on our website. Seek appropriate training to safely handle this product under normal conditions. Use the recommended personal protective equipment to prevent chemical exposures.

Bertin Bioreagent does not make any other warranty or representation whatsoever whether expressed or implied, with respect to these products. In no event will Bertin Bioreagent be liable for incidental, consequential or punitive damages.

Contact Bertin Bioreagent

Parc d'Activités du Pas du Lac  
10 bis avenue Ampère  
71810 Montigny le Bretonneux - France

Tel.: +33 (0)139 306 036

<https://www.bertin-bioreagent.com/pa206/contact-us>

<https://www.bertin-bioreagent.com>